Bangladesh



Monitoring the situation of children and women

Multiple Indicator Cluster Survey 2012-2013

Progotir Pathey



Key Findings



Government of the People's Republic of Bangladesh



Bangladesh Bureau of Statistics Statistics and Informatics Division (SID) Ministry of Planning



United Nations Children's Fund





Bangladesh

Multiple Indicator Cluster Survey 2012-2013

Progotir Pathey

Key Findings

May, 2014







The Bangladesh Multiple Indicator Cluster Survey (MICS) Progotir Pathey was carried out in 2012-2013 by Bangladesh Bureau of Statistics in collaboration with The United Nations Children's Fund (**UNICEF**), as part of the global MICS programme. Technical and financial support was provided by the United Nations Children's Fund (UNICEF).

The global MICS programme was developed by UNICEF in the 1990s as an international household survey programme to collect internationally comparable data on a wide range of indicators on the situation of children and women. MICS surveys measure key indicators that allow countries to generate data for use in policies and programmes, and to monitor progress towards the Millennium Development Goals (MDGs) and other internationally agreed upon commitments.

The objective of this report is to facilitate the timely dissemination and use of results from the Progotir Pathey MICS prior to the release of full tables and the final report that will contain detailed information on all survey findings by various demographic, social, economic and cultural characteristics. The final survey report is expected to be released in June 2014.

Results presented in this report are not expected to change and are considered final unless otherwise mentioned. For more information on indicators and the analysis conducted in the full final report please go to bbs.gov.bd, mics.unicef.org and childinfo.org.

Reprint: March 2015 Suggested citation:

Progotir Pathey Multiple Indicator Cluster Survey 2012-2013, Key Findings.

Bangladesh Bureau of Statistics and UNICEF Bangladesh 2014, Dhaka, Bangladesh

Contents

MICS AT A GLANCE	1
CHILD MORTALITY	2
Nutrition	3
CHILD HEALTH	9
WATER AND SANITATION	11
REPRODUCTIVE HEALTH	13
CHILD DEVELOPMENT	18
LITERACY AND EDUCATION	20
CHILD PROTECTION	23
HIV/AIDS	27
ACCESS TO MASS MEDIA AND ICT	29
Notes	30



BANGLADESH 2012-2013 MICS AT A GLANCE

Survey implemen	tation	Bangla	desh MICS
Sample frame	Population and Housing	Questionnaires	Household
	Census, 2011		Women (age 15-49)
			Children under five
		V	Vater quality testing
Training of Trainers	October 1-14, 2012	Fieldwork December	er 2012 – April 2013
Interviewer training	November 1-14, 2012		
Survey sample			
Households		Children under five	
Sampled	55,120	Eligible	23,402
Occupied	52,711	Mothers/caretakers interviewed	20,903
Interviewed	51,895	Response rate (Per cent)	89.3
Response rate (Per cent)	98.5		
Women			
Eligible for interviews	59,599		
Interviewed	51,791		
Response rate (Per cent)	86.9		

Survey population		Banglad	desh MICS
Average household size	4.6	Percentage of population living in	
		Urban areas	20.7
Percentage of population under:	0.0	Rural areas	79.3
Age 5	9.9	Division	
Age 18	39.0	Barisal	6.3
Development of warmen and 15 40 warms with at		Chittagong	20.1
Percentage of women age 15-49 years with at	45.2	Dhaka	30.7
least one live birth in the last 2 years	15.3	Khulna	11.2
		Rajshahi	13.0
		Rangpur	11.9
		Svlhet	6.7

Housing characteristics	
Percentage of households with	
Electricity	61.5
Finished floor	25.7
Finished roofing	97.4
Finished walls	29.4
Mean number of persons per room used for sleeping	2.67

Household or personal assets			
Percentage of households that own			
A television	37.7		
A refrigerator	14.2		
Electric fan	53.0		
Almirah / Wardrobe	41.4		
Agricultural land	43.0		
Farm animals/livestock	62.9		
Percentage of households where at leas	st a		
member has or owns a			
Watch	33.0		
Mobile phone	85.9		
Bicycle	28.7		
Computer	3.4		
Car or truck	0.7		




CHILD MORTALITY

Early childhood mortality Bangladesh MIG			MICS	
MIC	S cator	Indicator	Description	Value ^A
1.2	MDG 4.2	Infant mortality rate	Probability of dying between birth and the first birthday	
			National	46
			Division	
			Barisal	38
			Chittagong	36
			Dhaka	48
			Khulna	33
			Rajshahi	58
			Rangpur	49
			Sylhet	59
1.5	MDG 4.1	Under-five mortality rate	Probability of dying between birth and the fifth birthday	
			National	58
			Division	
			Barisal	47
			Chittagong	45
			Dhaka	62
			Khulna	41
			Rajshahi	78
			Rangpur	63
			Sylhet for to 2008 3. The West Model Life table was assumed to approxim	80

^AIndicator values as per 1,000 live births and refer to 2008.3. The West Model Life table was assumed to approximate the age pattern of mortality in Bangladesh and calculations are based on the time since first birth group version of the indirect children ever born/children surviving method for 5-9 years group.

Figure 1: Early childhood mortality rates, Bangladesh MICS, 2012-2013



Note: Indicator values are per 1,000 live births



NUTRITION


Indicator	Nutri	Nutritional status ⁱ Bangladesh MICS			
(a) Moderate and severe of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.1b Underweight prevalence (b) Severe of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Rhulna Ranghur Sylhet Savere of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Rhulna Rajshahi Rangpur Sylhet Savere of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Rhulna Rajshahi Rangpur Sylhet Savere of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Rhulna Rajshahi Rangpur Sylhet Savere of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Rajshahi Rangpur Sylhet Savere of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Rajshahi Rangpur Sylhet Savere of the median height for age of the WHO standard National Division Substantial Rangpur Sylhet Savere of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Chittagong Dhaka Chittagong Dhaka Chittagong Dhaka		or	Indicator	Description	Value
National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.1b Underweight prevalence (b) Severe (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe (c) Severe (c) Severe (d) Severe (d) Severe (e) Severe (e) Stunting prevalence (f) Severe (h) S	2.1a N	/IDG 1.8		minus two standard deviations (moderate and severe)	
Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.1b Underweight prevalence (b) Severe (b) Severe (b) Severe (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe (b) Severe (c) Stunting prevalence (b) Severe (c) Stunting prevalence (c) Severe (d) Severe (e) Stunting prevalence (b) Severe (c) Severe (d) Severe (d) Severe (e) Stunting prevalence (b) Severe (d) Severe (e) Stunting prevalence (e) Severe (f) Severe (h) Se				National	31.9
Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.1b Underweight prevalence (b) Severe Underweight prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Stunting prevalence (c) Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Of the median height for age of the WHO standard National Division Barisal Chittagong Division Barisal Chittagong Dhaka					35.2
Dhaka Khulna Rajshahi Rangpur Sylhet 2.1b Underweight prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe Of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe (c) Severe Of the median height for age of the WHO standard National Division Barisal Chittagong Of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka					32.2
Khulna Rajshahi Rangpur Sylhet 2.1b Underweight prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet Autional Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Akhulna Rajshahi Rangpur Sylhet Stunting prevalence (b) Severe Of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka					30.8
Rajshahi Rangpur Sylhet 2.1b Underweight prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet Percentage of children under age 5 who fall below minus two standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Of the median height for age of the WHO standard National Division Barisal Chittagong Of the median height for age of the WHO standard National Division Barisal Chittagong Of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Khulna	28.5
Rangpur Sylhet 2.1b Underweight prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Chittagong Division Barisal Chittagong Division					29.9
Sylhet 2.1b Underweight prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka All Division Barisal Chittagong Dhaka Division Barisal Chittagong Dhaka					32.6
2.1b Underweight prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median weight for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka					39.7
National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 5. 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka	2.1b			Percentage of children under age 5 who fall below	
Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				of the median weight for age of the WHO standard	
Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe (b) Severe Barisal Chittagong Dhaka				National	8.8
Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe Fercentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Division	
Dhaka Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka 1 Chittagong Dhaka				Barisal	9.7
Khulna Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (b) Severe Stunting prevalence (a) Moderate and severe (b) Severe Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Stunting prevalence (b) Severe Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Stunting prevalence (c) Severe Stunting prevalence (d) Severe Stunting prevalence (e) Severe Stunting prevalence (f) Severe Stunting prevalence (h) Severe (h) Se				Chittagong	9.2
Rajshahi Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (b) Severe Rajshahi Rangpur Sylhet Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Dhaka	8.9
Rangpur Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Chittagong Dhaka				Khulna	6.2
Sylhet 2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (a) Moderate and severe (b) of the median height for age of the WHO standard (b) National (b) Division (c) Dhaka (c) Dhak				Rajshahi	7.2
2.2a Stunting prevalence (a) Moderate and severe (a) Moderate and severe Percentage of children under age 5 who fall below minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka 1 Chittagong Dhaka					8.1
(a) Moderate and severe minus two standard deviations (moderate and severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Sylhet	13.1
of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka	2.2a		Stunting prevalence	Percentage of children under age 5 who fall below	
Division Barisal Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka			(a) Moderate and severe		
Chittagong Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka					42.0
Dhaka Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below (b) Severe minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Barisal	41.4
Khulna Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below (b) Severe minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Chittagong	43.1
Rajshahi Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below (in the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Dhaka	42.1
Rangpur Sylhet 2.2b Stunting prevalence (b) Severe Percentage of children under age 5 who fall below minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Khulna	34.4
Sylhet 2.2b Stunting prevalence Percentage of children under age 5 who fall below (b) Severe minus three standard deviations (severe) of the median height for age of the WHO standard National Division Barisal Chittagong Dhaka				Rajshahi	39.4
2.2b Stunting prevalence Percentage of children under age 5 who fall below (b) Severe minus three standard deviations (severe) of the median height for age of the WHO standard National 1 Division Barisal 1 Chittagong 1 Dhaka				Rangpur	43.7
(b) Severe minus three standard deviations (severe) of the median height for age of the WHO standard National 1 Division Barisal 1 Chittagong 1 Dhaka				Sylhet	50.6
National 1 Division Barisal 1 Chittagong 1 Dhaka 1	2.2b		• • • • • • • • • • • • • • • • • • • •	minus three standard deviations (severe)	
Barisal 1 Chittagong 1 Dhaka 1				National	16.4
Chittagong 1 Dhaka 1					15.8
Dhaka 1]				18.7
					16.7
Nituria 1]				11.3
Rajshahi 1					13.2
·				•	16.2
					22.2



Nutritional status ⁱ Bangladesh			MICS
MICS Indicator	Indicator	Description	Value
2.3a	Wasting prevalence	Percentage of children under age 5 who fall below	
	(a) Moderate and severe	minus two standard deviations (moderate and severe)	
		of the median weight for height of the WHO standard	
		National	9.6
		Division	
		Barisal	11.7
		Chittagong	9.2
		Dhaka	9.2
		Khulna	10.0
		Rajshahi	9.1
		Rangpur	8.7
		Sylhet	13.3
2.3b	Wasting prevalence	Percentage of children under age 5 who fall below	
	(b) Severe	minus three standard deviations (severe)	
		of the median weight for height of the WHO standard	
		National	1.6
		Division	
		Barisal	1.5
		Chittagong	1.8
		Dhaka	1.3
		Khulna	1.7
		Rajshahi	1.5
		Rangpur	1.4
		Sylhet	2.8
2.4	Overweight prevalence	Percentage of children under age 5 who are above two	
		standard deviations of the median weight for height of the	
		WHO standard	
		National	1.6
		Division	
		Barisal	0.8
		Chittagong	1.2
		Dhaka	2.2
		Khulna	1.1
		Rajshahi	1.2
		Rangpur	1.1
		Sylhet	2.9



Figure 2: Underweight, stunted, wasted and overweight children under age 5 (moderate and severe), Bangladesh MICS, 2012-2013





	eding and infant f	eeding Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
2.5	Children ever breastfed	Percentage of women with a live birth in the last 2 years who breastfed their last live-born child at any time	
		National	97.1
		Division	
		Barisal	96.4
		Chittagong	95.0
		Dhaka	97.8
		Khulna	97.6
		Rajshahi	97.8
		Rangpur	98.7
		Sylhet	97.8
2.6	Early initiation of	Percentage of women with a live birth in the last 2 years	
	breastfeeding	who put their last newborn to the breast within one hour of	
		birth	
		National	57.4
		Division	
		Barisal	55.8
		Chittagong	53.1
		Dhaka	59.6
		Khulna	47.3
		Rajshahi	56.5
		Rangpur	59.1
		Sylhet	73.5
2.7	Exclusive breastfeeding	Percentage of infants under 6 months of age who are	
	under 6 months	exclusively breastfed ⁱⁱ	
		National	56.4
		Division	
		Barisal	65.0
		Chittagong	69.4
		Dhaka	46.2
		Khulna	50.5
		Rajshahi	50.1
		Rangpur	67.7
		Sylhet	52.2
2.8	Predominant breastfeeding	Percentage of infants under 6 months of age who received	
	under 6 months	breast milk as the predominant source of nourishmentiii	
		during the previous day	
		National	71.9
		Division	
		Barisal	74.5
		Chittagong	82.8
		Dhaka	64.8
		Khulna	73.8
		Rajshahi	61.0
		Rangpur	75.7
		Sylhet	73.9



Breastfe	eding and infant f	feeding Bangladesh	n MICS
MICS Indicator	Indicator	Description	Value
2.9	Continued breastfeeding at 1 year	Percentage of children age 12-15 months who received breast milk during the previous day	
	·	National	95.3
		Division	
		Barisal	94.3
		Chittagong	93.2
		Dhaka	97.3
		Khulna	96.3
		Rajshahi	92.7
		Rangpur	98.2
		Sylhet	94.3
2.10	Continued breastfeeding at	Percentage of children age 20-23 months who received	54.5
2.10	2 years	breast milk during the previous day	
	2 years	National	07.5
			87.5
		Division	00.5
		Barisal	90.5
		Chittagong	77.7
		Dhaka	91.0
		Khulna	92.4
		Rajshahi	91.9
		Rangpur	92.8
		Sylhet	85.5
2.11	Median duration of	The age in months when 50 percent of children age 0-35	
	breastfeeding	months did not receive breast milk during the previous day	
		National	32.1
		Division	
		Barisal	33.3
		Chittagong	25.1
		Dhaka	33.0
		Khulna	≥36.0
		Rajshahi	≥36.0
		Rangpur	33.9
		Sylhet	29.7
2.12	Age-appropriate	Percentage of children age 0-23 months appropriately	
	breastfeeding	fed ^{iv} during the previous day	
		National	66.5
		Division	
		Barisal	62.0
		Chittagong	58.3
		Dhaka	65.5
		Khulna	76.2
		Rajshahi	70.2
		Rangpur	76.8
		Sylhet	63.0
2.13	Introduction of solid som:	Percentage of infants age 6-8 months who received solid,	03.0
2.13	Introduction of solid, semi-	semi-solid or soft foods during the previous day	42.4
	solid or soft foods	semi-sond or soft foods during the previous day	42.4



Breastfeeding and infant feeding Bangladesh			h MICS
MICS Indicator	Indicator	Description	Value
2.18	Bottle feeding	Percentage of children age 0-23 months who were fed with	
		a bottle during the previous day	
		National	12.1
		Division	
		Barisal	10.2
		Chittagong	11.1
		Dhaka	17.4
		Khulna	8.6
		Rajshahi	15.5
		Rangpur	5.4
		Sylhet	4.5

Salt iodization		Bangladesh N	
MICS Indicator	Indicator	Description	Value
2.19	lodized salt consumption	Percentage of households with salt testing 15 parts per million or more of iodate	
		National	54.3
		Division	
		Barisal	62.4
		Chittagong	59.3
		Dhaka	64.6
		Khulna	60.1
		Rajshahi	36.3
		Rangpur	33.8
		Sylhet	50.7

Low-birt	th weight	Bangladesh MICS	
MICS Indicator	Indicator	Description	Value
2.20	Low-birthweight infants	Percentage of most recent live births in the last 2 years weighing below 2,500 grams at birth	
		National	26.0
		Division	
		Barisal	26.3
		Chittagong	29.2
		Dhaka	25.0
		Khulna	24.2
		Rajshahi	22.6
		Rangpur	24.2
		Sylhet	29.3
2.21	Infants weighed at birth	Percentage of most recent live births in the last 2 years who	
		were weighed at birth	
		National	35.9
		Division	
		Barisal	17.5
		Chittagong	25.8
		Dhaka	40.6
		Khulna	44.3
		Rajshahi	34.5
		Rangpur	59.3
		Sylhet	19.5



CHILD HEALTH

Tetanus to	Bangladesh	MICS	
MICS Indicator	Indicator	Description	Value
3.9	Neonatal tetanus protection	Percentage of women age 15-49 years with a live birth in the last 2 years who were given at least two doses of tetanus toxoid vaccine within the appropriate interval prior to the most recent birth	
		National Division	80.8
		Barisal Chittagong	79.4 79.4
		Dhaka Khulna	82.5 83.4
		Rajshahi Rangpur	85.0 83.4
		Sylhet	66.7

Diarrhoea		Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
-	Children with diarrhoea	Percentage of children under age 5 with diarrhoea in the last	
		2 weeks	
		National	3.9
		Division	
		Barisal	6.3
		Chittagong	4.5
		Dhaka	3.5
		Khulna	3.3
		Rajshahi	3.5
		Rangpur	3.8
		Sylhet	3.8
3.12	Diarrhoea treatment with	Percentage of children under age 5 with diarrhoea in the last	
	oral rehydration therapy	2 weeks who received ORT (ORS packet, pre-packaged ORS	
	(ORT) and continued	fluid, recommended homemade fluid or increased fluids)	
	feeding	and continued feeding during the episode of diarrhoea	
		National	64.6
		Division	
		Barisal	59.5
		Chittagong	68.7
		Dhaka	64.2
		Khulna	65.1
		Rajshahi	65.2
		Rangpur	67.6
		Sylhet	51.6



Acute Re	spiratory Infection	n (ARI) symptoms Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
	Children with ARI	Percentage of children under age 5 with ARI symptoms in	
	symptoms	the last 2 weeks	
		National	3.2
		Division	
		Barisal	4.1
		Chittagong	3.6
		Dhaka	2.1
		Khulna	4.3
		Rajshahi	3.1
		Rangpur	3.2
		Sylhet	4.7
3.13	Care-seeking for children	Percentage of children under age 5 with ARI symptoms in	
	with ARI symptoms	the last 2 weeks for whom advice or treatment was sought	
		from a health facility or provider	
		National	35.8
		Division	
		Barisal	26.1
		Chittagong	30.0
		Dhaka	37.8
		Khulna	46.2
		Rajshahi	35.0
		Rangpur	37.7
		Sylhet	39.2
3.14	Antibiotic treatment for	Percentage of children under age 5 with ARI symptoms in	
	children with ARI	the last 2 weeks who received antibiotics	
	symptoms		
		National	74.3
		Division	
		Barisal	78.4
		Chittagong	66.8
		Dhaka	82.3
		Khulna	76.4
		Rajshahi	69.4
		Rangpur	72.8
		Sylhet	78.6

Solid fue	l use	Bangladesh MICS	
MICS Indicator	Indicator	Description	Value
3.15	Use of solid fuels for	Percentage of household members in households that use	
	cooking	solid fuels as the primary source of domestic energy to cook	
		National	88.2
		Division	
		Barisal	98.3
		Chittagong	91.1
		Dhaka	75.9
		Khulna	95.9
		Rajshahi	92.9
		Rangpur	95.9
		Sylhet	91.1




WATER AND SANITATION

Wat	er and	d sanitation	Bangladesh	MICS
MICS Indica	tor	Indicator	Description	Value
4.1	MDG 7.8	Use of improved drinking	Percentage of household members using improved sources	
		water sources	of drinking water	
			National	97.9
			Division	
			Barisal	95.3
			Chittagong	97.0
			Dhaka	99.9
			Khulna	94.4
			Rajshahi	99.3
			Rangpur	99.9
			Sylhet	93.8
4.2		Water treatment	Percentage of household members in households using	
			unimproved drinking water who use an appropriate	
			treatment method	
			National	25.6
			Division	
			Barisal	45.1
			Chittagong	5.1
			Dhaka	16.6
			Khulna	40.7
			Rajshahi	0.0
			Rangpur	0.0
			Sylhet	26.2
4.3	MDG 7.9	Use of improved sanitation	Percentage of household members using improved	
			sanitation facilities which are not shared	
			National	55.9
			Division	
			Barisal	52.0
			Chittagong	59.4
			Dhaka	54.0
			Khulna	58.0
			Rajshahi	52.0
			Rangpur	57.4
			Sylhet	58.6
4.4		Safe disposal of child's	Percentage of children age 0-2 years whose last stools were	
		faeces	disposed of safely	
			National	38.7
			Division	
			Barisal	39.3
			Chittagong	37.6
			Dhaka	46.0
			Khulna	43.8
			Rajshahi	34.6
			Rangpur	21.4
			Sylhet	35.5



Water ar	nd sanitation	Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
4.5	Place for handwashing	Percentage of households with a specific place for hand washing where water and soap or other cleansing agent are present	
		National Division	59.1
		Barisal	50.0
		Chittagong	71.4
		Dhaka	50.1
		Khulna	66.5
		Rajshahi	67.9
		Rangpur	61.5
		Sylhet	54.2
4.6	Availability of soap or other	Percentage of households with soap or other cleansing	
	cleansing agent	agent	
		National	94.0
		Division	
		Barisal	94.9
		Chittagong	93.1
		Dhaka	93.8
		Khulna	95.3
		Rajshahi	93.5
		Rangpur	95.6
		Sylhet	92.5

Figure 4: Use of improved water and sanitation in urban and rural areas, Bangladesh MICS, 2012-2013



Percentage of household members using an improved water source

Percentage of household members using an improved sanitation facility which is not shared



REPRODUCTIVE HEALTH

	ility, co	ontraception and	l unmet need Bangladesh	MICS
MICS Indica	itor	Indicator	Description	Value
		Total fertility rate	Total fertility rate ^A for women age 15-49 years	
			National	2.3
			Division	
			Barisal	2.3
			Chittagong	2.7
			Dhaka	2.3
			Khulna	1.9
			Rajshahi	1.9
			Rangpur	2.2
			Sylhet	2.9
5.1	MDG 5.4	Adolescent birth rate	Age-specific fertility rate ^A for women age 15-19 years	
			National	83
			Division	
			Barisal	75
			Chittagong	86
			Dhaka	82
			Khulna	84
			Rajshahi	99
			Rangpur	94
			Sylhet	45
5.2		Early childbearing	Percentage of women age 20-24 years who had at least one	
			live birth before age 18	
			National	24.4
			Division	
			Barisal	21.3
			Chittagong	19.7
			Dhaka	23.3
			Khulna	27.1
			Rajshahi	34.0
			Rangpur	30.5
			Sylhet	14.8
5.3	MDG 5.3	Contraceptive prevalence	Percentage of women age 15-49 years currently married	
		rate	who are using (or whose partner is using) a (modern or	
			traditional) contraceptive method	
			National	61.8
			Division	
			Barisal	56.8
			Chittagong	53.0
			Dhaka	60.1
			Khulna	70.3
			Rajshahi	68.1
			Rangpur	72.9
			Sylhet	46.5




Fer	tility, co	ontraception	and unmet need	Bangladesh MICS
MICS		Indicator	Description	Value
5.4	MDG 5.6	Unmet need	Percentage of women age 15-49 married or in union who are fect births or limit the number of chinot currently using contraceptic	und and want to space their ildren they have and who are
			National Division	13.9
			Barisal	19.0
			Chittagong	18.9
			Dhaka	15.0
			Khulna	9.3
			Rajshahi	10.2
			Rangpur	9.1
			Sylhet	16.3

^AThe age-specific fertility rate is defined as the number of live births to women in a specific age group during a specified period, divided by the average number of women in that age group during the same period, expressed per 1,000 women. The age-specific fertility rate for women age 15-19 years is also termed as the adolescent birth rate.

The total fertility rate (TFR) is calculated by summing the age-specific fertility rates calculated for each of the 5-year age groups of women, from age 15 through to age 49. The TFR denotes the average number of children to which a woman will have given birth by the end of her reproductive years (by age 50) if current fertility rates prevailed.

Figure 5: Use of contraception, Bangladesh MICS, 2012-2013





Maternal and newborn health Bangladesh M			MICS	
MICS		Indicator	Description	Value
5.5a		Antenatal care coverage	Percentage of women age 15-49 years with a live birth in the last 2 years who were attended during their last pregnancy that led to a live birth at least once by skilled health personnel	
			National Division	58.7
			Barisal	40.3
			Chittagong	58.1
			Dhaka	61.9
			Khulna	74.6
			Rajshahi	63.6
			Rangpur	46.6
			Sylhet	52.1
5.5b	MDG 5.5	Antenatal care coverage	Percentage of women age 15-49 years with a live birth in the last 2 years who were attended during their last pregnancy that led to a live birth at least four times by any	32
			provider	24:
			National	24.7
			Division	444
			Barisal	14.0
			Chittagong	21.
			Dhaka Wasta a	26.3
			Khulna Baishahi	27.:
			Rajshahi	25.6
			Rangpur Sylhet	35.8 16.0
5.6		Content of antenatal care	Percentage of women age 15-49 years with a live birth in the last 2 years who had their blood pressure measured and	10.0
			gave urine and blood samples during the last pregnancy that led to a live birth	
			National	38.0
			Division	
			Barisal	28.4
			Chittagong	36.
			Dhaka	44.3
			Khulna	45.5
			Rajshahi	31.3
			Rangpur	28.2
			Sylhet	37.6
5.7	MDG 5.2	Skilled attendant at delivery	Percentage of women age 15-49 years with a live birth in the last 2 years who were attended by skilled health personnel	
		,	during their most recent live birth	
			National Division	43.5
			Barisal	38.4
			Chittagong	41.5
			Dhaka	44.8
			Khulna	56.7
			Rajshahi	51.8
			Rangpur	39.4
			Sylhet	26.7




Materna	Maternal and newborn health Bangladesh		MICS
MICS Indicator	Indicator	Description	Value
5.8	Institutional deliveries	Percentage of women age 15-49 years with a live birth in the last 2 years whose most recent live birth was delivered in a health facility	
		National Division	31.0
		Barisal Chittagong	17.1 27.1
		Dhaka Khulna	34.9
		Rajshahi	45.6 38.1
		Rangpur Sylhet	23.0 20.8
5.9	Caesarean section	Percentage of women age 15-49 years whose most recent live birth in the last 2 years was delivered by caesarean section	
		National Division	19.1
		Barisal	10.5
		Chittagong	14.5
		Dhaka	24.4
		Khulna	30.5
		Rajshahi	22.4
		Rangpur	11.7
		Sylhet	10.8

Post-natal health checks Bangladesh I			MICS
MICS Indicator	Indicator	Description	Value
5.10	Post-partum stay in health facility	Percentage of women age 15-49 years who stayed in the health facility for 12 hours or more after the delivery of their most recent live birth in the last 2 years	
		National Division	82.8
		Barisal	85.9
ı		Chittagong	80.4
		Dhaka	86.8
		Khulna	83.5
		Rajshahi	80.4
		Rangpur	74.2
		Sylhet	81.0
5.11	Post-natal health check for the newborn	Percentage of last live births in the last 2 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery	
		National	41.2
		Division	
		Barisal	22.0
		Chittagong	33.6
		Dhaka	43.9
		Khulna	52.0
		Rajshahi	41.9
		Rangpur	46.8
		Sylhet	45.3



Post-natal health checks		Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
5.12	Post-natal health check for the mother	Percentage of women age 15-49 years who received a health check while in facility or at home following delivery, or a post-natal care visit within 2 days after delivery of their most recent live birth in the last 2 years	
		National Division	40.4
		Barisal	23.3
		Chittagong	34.3
		Dhaka	43.3
		Khulna	50.4
		Rajshahi	39.3
		Rangpur	43.2
		Sylhet	45.3

Figure 6: Place of delivery and post-natal health checks, Bangladesh MICS, 2012-2013





CHILD DEVELOPMENT

Child dev	velopment	Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
6.1	Attendance to early childhood education	Percentage of children age 36-59 months who are attending an early childhood education programme	
		National	13.4
		Division	
		Barisal	18.4
		Chittagong	11.7
		Dhaka	15.6
		Khulna	13.6
		Rajshahi	10.2
		Rangpur	13.2
		Sylhet	10.5
6.2	Support for learning	Percentage of children age 36-59 months with whom an	
		adult has engaged in four or more activities to promote	
		learning and school readiness in the last 3 days	
		National ,	78.0
		Division	
		Barisal	80.4
		Chittagong	76.5
		Dhaka	78.9
		Khulna	85.9
		Rajshahi	75.1
		Rangpur	75.0
		Sylhet	76.4
6.3	Father's support for	Percentage of children age 36-59 months whose biological	, , , ,
0.0	learning	father has engaged in four or more activities to promote	
		learning and school readiness in the last 3 days	
		National	10.1
		Division	
		Barisal	4.0
		Chittagong	9.1
		Dhaka	10.2
		Khulna	8.7
		Rajshahi	6.3
		Rangpur	20.6
		Sylhet	9.0
6.4	Mother's support for	Percentage of children age 36-59 months whose biological	
	learning	mother has engaged in four or more activities to promote	
	5	learning and school readiness in the last 3 days	
		National	40.8
		Division	
		Barisal	28.5
		Chittagong	39.5
		Dhaka	44.1
		Khulna	41.5
		Rajshahi	41.9
		Rangpur	45.9
		Manippan	30.7



Child dev	velopment	Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
6.5	Availability of children's books	Percentage of children under age 5 who have three or more children's books	
		National	8.8
		Division	
		Barisal	9.6
		Chittagong	7.9
		Dhaka	11.5
		Khulna	10.3
		Rajshahi	8.7
		Rangpur	5.4
		Sylhet	3.8
6.6	Availability of playthings	Percentage of children under age 5 who play with two or	
		more types of playthings	60.0
		National	60.3
		Division	60.0
		Barisal	60.9
		Chittagong	44.7
		Dhaka	57.6
		Khulna	72.8
		Rajshahi	69.8
		Rangpur	73.5
		Sylhet	68.1
6.7	Inadequate care	Percentage of children under age 5 left alone or in the care	
		of another child younger than 10 years of age for more than one hour at least once in the last week	
			11.0
		National	11.6
		Division	10.2
		Barisal	10.3
		Chittagong Dhaka	15.7
		Khulna	10.2
			7.5 10.7
		Rajshahi	14.3
		Rangpur Sylhet	8.3
6.8	Early child development	Percentage of children age 36-59 months who are	0.3
0.6	index	developmentally on track in at least three of the following	
	ilidex	four domains: literacy-numeracy, physical, social-emotional,	
		and learning	
		National	63.9
		Division	03.3
		Barisal	67.3
		Chittagong	54.4
		Dhaka	65.1
		Khulna	69.0
		Rajshahi	65.2
		Rangpur	77.7

LITERACY AND EDUCATION

Lite	racy an	d education ^v	Bangladesh I	MICS
MICS	Indicator	Indicator	Description	Value
7.1	MDG	Literacy rate among young	Percentage of young women age 15-24 years who are able	
	2.3	people	to read a short simple statement about everyday life or who	
			attended secondary or higher education	
			National	82.0
			Division	
			Barisal	86.1
			Chittagong	81.5
			Dhaka	80.9
			Khulna	87.8
			Rajshahi	83.0
			Rangpur	80.0
			Sylhet	78.2
7.2		School readiness	Percentage of children in first grade of primary school who	
			attended pre-school during the previous school year	
			National	43.5
			Division	
			Barisal	76.4
			Chittagong	35.9
			Dhaka	49.9
			Khulna	52.2
			Rajshahi	30.2
			Rangpur	36.2
			Sylhet	29.2
7.3		Net intake rate in primary	Percentage of children of school-entry age who enter the	
		education	first grade of primary school	
			National	33.1
			Division	
			Barisal	32.3
			Chittagong	35.8
			Dhaka	33.1
			Khulna	27.5
			Rajshahi	32.3
			Rangpur	41.0
			Sylhet	23.1
7.4	MDG	Primary school net	Percentage of children of primary school age currently	
	2.1	attendance ratio (adjusted)	attending primary or secondary school	
			National	73.2
			Division	
			Barisal	71.9
			Chittagong	72.4
			Dhaka	72.5
			Khulna	75.4
			Rajshahi	75.1
			Rangpur	75.7
			Sylhet	69.4




		d education ^v	Bangladesh	
	ndicator	Indicator	Description	Value
7.5		Secondary school net	Percentage of children of secondary school age currently	
		attendance ratio (adjusted)	attending secondary school or higher	
			National	46.1
			Division	
			Barisal	47.8
			Chittagong	45.6
			Dhaka	45.6
			Khulna	50.9
			Rajshahi	46.1
			Rangpur	52.3
			Sylhet	32.7
7.6	MDG 2.2	Children reaching last grade of primary	Percentage of children entering the first grade of primary school who eventually reach last grade	
			National	96.4
			Division	
			Barisal	96.5
			Chittagong	96.9
			Dhaka	96.1
			Khulna	96.7
			Rajshahi	96.9
			Rangpur	97.5
			Sylhet	93.4
7.7		Primary completion rate	Number of children attending the last grade of primary	
			school (excluding repeaters) divided by number of children	
			of primary school completion age (age appropriate to final	
			grade of primary school)	
			National	79.5
			Division	
			Barisal	79.5
			Chittagong	80.8
			Dhaka	71.2
			Khulna	82.4
			Rajshahi	92.5
			Rangpur	87.4
			Sylhet	73.5
7.8		Transition rate to	Number of children attending the last grade of primary	
		secondary school	school during the previous school year who are in the first	
			grade of secondary school during the current school year	
			divided by number of children attending the last grade of	
			primary school during the previous school year	
			National	94.7
			Division	
			Barisal	95.8
			Chittagong	94.7
			Dhaka	94.5
			Khulna	96.8
			Rajshahi	93.8
			Rangpur	94.4
			Sylhet	93.2

Literacy and education ^v Bar		Bangladesh	MICS	
MICS	Indicator	Indicator	Description	Value
7.9	MDG	Gender parity index	Primary school net attendance ratio (adjusted) for girls	
	3.1	(primary school)	divided by primary school net attendance ratio (adjusted)	
			for boys	
			National	1.07
			Division	
			Barisal	1.10
			Chittagong	1.07
			Dhaka	1.07
			Khulna	1.08
			Rajshahi	1.06
			Rangpur	1.04
			Sylhet	1.10
7.10	MDG	Gender parity index	Secondary school net attendance ratio (adjusted) for girls	
	3.1	(secondary school)	divided by secondary school net attendance ratio (adjusted)	
			for boys	
			National	1.30
			Division	
			Barisal	1.35
			Chittagong	1.51
			Dhaka	1.24
			Khulna	1.26
			Rajshahi	1.23
			Rangpur	1.25
			Sylhet	1.25

NOTE: Field data was collected over a period spanning two school years 2012 and 2013. Though interviewers were instructed to retain the same reference period for education questions, different reference periods seem to have been used in some cases. The data here refers to the age at the beginning of the school year 2012 only. The education dataset is under review and results may be subject to change.

Figure 7: Education indicators by sex, Bangladesh MICS, 2012-2013





CHILD PROTECTION

Birth registration		Bangladesh	
MICS Indicator	Indicator	Description	Value
8.1	Birth registration	Percentage of children under age 5 whose births are reported registered	
		National	37.0
		Division	
		Barisal	32.3
		Chittagong	41.4
		Dhaka	34.3
		Khulna	32.1
		Rajshahi	32.6
		Rangpur	47.6
		Sylhet	35.0

Child discipline		Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
8.3	Violent discipline	Percentage of children age 1-14 years who experienced	
		psychological aggression or physical punishment during the	
		last one month	
		National	82.3
		Division	
		Barisal	88.6
		Chittagong	81.3
		Dhaka	81.3
		Khulna	86.9
		Rajshahi	80.2
		Rangpur	79.5
		Sylhet	85.3

Figure 8: Child disciplining methods, children age 1-14 years, Bangladesh MICS, 2012-2013



Early ma	Early marriage and polygyny Banglades		sh MICS	
MICS Indicator	Indicator	Description	Value	
8.4	Marriage before age 15	Percentage of women age 15-49 years who were first married or in union before age 15		
		National	23.8	
		Division		
		Barisal	20.8	
		Chittagong	14.5	
		Dhaka	23.1	
		Khulna	31.1	
		Rajshahi	33.3	
		Rangpur	31.5	
		Sylhet	9.2	
8.5	Marriage before age 18	Percentage of women age 20-49 years who were first married or in union before age 18		
		National	62.8	
		Division	02.8	
		Barisal	65.8	
			50.5	
		Chittagong Dhaka	61.4	
		Khulna	71.1	
			71.1	
		Rajshahi	72.8 75.9	
		Rangpur Sylhet	75.9 38.5	
8.6	Young people age 15-19	Percentage of young women age 15-19 years who are		
	years currently married or in union	married or in union		
		National	34.3	
		Division		
		Barisal	31.7	
		Chittagong	27.7	
		Dhaka	33.3	
		Khulna	43.5	
		Rajshahi	47.8	
		Rangpur	41.9	
		Sylhet	13.7	
8.7	Polygyny	Percentage of women age 15-49 years who are in a		
		polygynous union		
		National	4.2	
		Division		
		Barisal	3.9	
		Chittagong	2.9	
		Dhaka	4.0	
		Khulna	4.1	
		Rajshahi	4.1	
		Rangpur	5.0	
		Sylhet	8.7	



Early marriage and polygyny Banglade		yny Bangladesh	MICS
MICS Indicator	Indicator	Description	Valu
8.10a	Spousal age difference	Percentage of young women who are married or in union and whose spouse is 10 or more years older among women age 15-19 years	
		National Division	20.
		Barisal	22.
		Chittagong	24.
		Dhaka	19
		Khulna	23
		Rajshahi	17
		Rangpur	15
		Sylhet	22
8.10b	Spousal age difference	Percentage of young women who are married or in union and whose spouse is 10 or more years older among women age 20-24 years	
		National	21
		Division	
		Barisal	17
		Chittagong	27
		Dhaka	19
		Khulna	23
		Rajshahi	19
		Rangpur	15
		Sylhet	32.

Children's	s living arrangeme	ents Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
8.13	Children's living arrangements	Percentage of children age 0-17 years living with neither biological parent	
		National	3.8
		Division	
		Barisal	4.5
		Chittagong	1.8
		Dhaka	4.2
		Khulna	4.5
		Rajshahi	5.3
		Rangpur	5.1
		Sylhet	2.4
8.14	Prevalence of children with	Percentage of children age 0-17 years with one or both	
	one or both parents dead	biological parents dead	
	·	National	4.3
		Division	
		Barisal	4.9
		Chittagong	4.9
		Dhaka	4.1
		Khulna	3.6
		Rajshahi	3.2
		Rangpur	4.1
		Sylhet	6.2
8.15	Children with at least one	Percentage of children 0-17 years with at least one	
	parent living abroad	biological parent living abroad	
		National	4.8
		Division	
		Barisal	2.3
		Chittagong	9.9
		Dhaka	5.0
		Khulna	2.8
		Rajshahi	1.5
		Rangpur	0.3
		Sylhet	6.3



HIV/AIDS

HIV	/AIDS kr	nowledge and att	itudes Bangladesh	MICS
MICS	Indicator	Indicator	Description	Value
-		Have heard of AIDS	Percentage of people age 15-49 years who have heard of	
			AIDS	
			National	55.8
			Division	
			Barisal	50.2
			Chittagong	58.1
			Dhaka	58.6
			Khulna	76.6
			Rajshahi	52.4
			Rangpur	38.3
			Sylhet	42.8
9.1	MDG 6.3	Knowledge about HIV	Percentage of young Women age 15-24 years who	
		prevention among young	correctly identify ways of preventing the sexual	
		people	transmission of HIV ^{vi} , and who reject major	
			misconceptions about HIV transmission	
			National	9.1
			Division	
			Barisal	7.4
			Chittagong	5.5
			Dhaka	13.0
			Khulna	9.4
			Rajshahi	10.0
			Rangpur	5.9
			Sylhet	5.3
9.2		Knowledge of mother-to-	Percentage of women age 15-49 years who correctly	
		child transmission of HIV	identify all three means ^{vii} of mother-to-child transmission of HIV	
			National	21.7
			Division	
			Barisal	26.1
			Chittagong	20.4
			Dhaka	19.8
			Khulna	32.9
			Rajshahi	25.2
			Rangpur	15.3
			Sylhet	14.0
9.3		Accepting attitudes	Percentage of women age 15-49 years expressing	
		towards people living with HIV	accepting attitudes on all four questionsviii toward people living with HIV	
			National	37.2
			Division	37.2
			Barisal	39.9
			Chittagong	41.4
			Dhaka	40.2
			Khulna	29.8
			Rajshahi	37.1
			Rangpur	33.8
			Sylhet	28.1

HIV testir	ng	Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
9.4	People who know where to be tested for HIV	Percentage of women age 15-49 years who state knowledge of a place to be tested for HIV	
		National	11.3
		Division	
		Barisal	8.2
		Chittagong	13.8
		Dhaka	11.0
		Khulna	16.4
		Rajshahi	10.5
		Rangpur	8.5
		Sylhet	6.2
9.7	HIV counselling	Percentage of women age 15-49 years who had a live birth in	
	during antenatal care	the last 2 years and received antenatal care during the	
		pregnancy of their most recent birth, reporting that they	
		received counselling on HIV during antenatal care	
		National	2.5
		Division	
		Barisal	1.4
		Chittagong	2.8
		Dhaka	2.6
		Khulna	2.7
		Rajshahi	2.2
		Rangpur	1.9
		Sylhet	2.8

Orp	hans		Bangladesh	MICS
MICS		Indicator	Description	Value
9.16	MDG 6.4	Ratio of school attendance of orphans to school attendance of non-orphans	Proportion attending school among children age 10-14 years who have lost both parents divided by proportion attending school among children age 10-14 years whose parents are alive and who are living with one or both parents	0.88



ACCESS TO MASS MEDIA AND ICT

Access to mass media		Bangladesh MI	
MICS Indicator	Indicator	Description	Value
10.1	10.1 Exposure to mass media Percentage of women age 15-49 years who, at least once a week, read a newspaper or magazine, listen to the radio, and watch television		
		National	1.6
		Division	
		Barisal	0.8
		Chittagong	0.6
		Dhaka	2.3
		Khulna	2.1
		Rajshahi	1.0
		Rangpur	0.9
		Sylhet	2.7

Use of in	nformation/com	munication technology Bangladesh	MICS
MICS Indicator	Indicator	Description	Value
10.2	Use of computers	Percentage of young women age 15-24 years who used a computer during the last 12 months	
		National	6.1
		Division Barisal	3.4
		Chittagong	4.9
		Dhaka Khulna	8.8 6.8
		Rajshahi	5.4
		Rangpur	3.3
		Sylhet	4.5
10.3	Use of internet	Percentage of young women age 15-24 years who used the internet during the last 12 months	
		National	3.0
		Division	
		Barisal	0.9
		Chittagong	2.1
		Dhaka	5.3
		Khulna	2.0
		Rajshahi	1.7
		Rangpur	1.4
		Sylhet	3.3

NOTES

ⁱWeight and height/length measurements were successfully completed for 95.3 and 92.9 per cent of children under age 5, respectively

ilnfants receiving breast milk, and not receiving any other fluids or foods, with the exception of oral rehydration solution, vitamins, mineral supplements and medicines

iilnfants receiving breast milk and certain fluids (water and water-based drinks, fruit juice, ritual fluids, oral rehydration solution, drops, vitamins, minerals, and medicines), but do not receive anything else (in particular, non-human milk and food-based fluids)

^{iv}Infants age 0-5 months who are exclusively breastfed, and children age 6-23 months who are breastfed and ate solid, semi-solid or soft foods

^vEducation indicators, wherever applicable, are based on information on reported school attendance (at any time during the school year), as a proxy for enrolment.

viUsing condoms and limiting sex to one faithful, uninfected partner

viiTransmission during pregnancy, during delivery, and by breastfeeding

viiiPeople (1) who think that a female teacher with the AIDS virus should be allowed to teach in school, (2) who would buy fresh vegetables from a shopkeeper or vendor who has the AIDS virus, (3) who would not want to keep it as a secret if a family member became infected with the AIDS virus, and (4) who would be willing to care for a family member who became sick with the AIDS virus